



Effects of a Natural Extract on Molecular Signature in Post-Acute Sequelae SARS-CoV-2 (PASC) Infection: A Quasi-Experimental Study

Alben Sigamani*, Sangeetha Sampath Kumar, K Sunil Naik

Carmel Research, Bangalore, India

ABSTRACT

Aim: Long COVID becomes an economic and public health challenge that affects the daily activities and quality of life of millions of COVID-19 survivors. Long COVID symptoms, particularly persistent fatigue, appear to be associated with a chronic state of inflammation. Resetting the inflammation and immune systems to baseline can reverse symptoms of Long COVID. The purpose of this study was to assess the efficacy and safety of CelWel in patients with long COVID.

Methods: This was a non-randomized, open-label pilot study with 15 COVID-19 infected male and female subjects who had Long COVID symptoms. Subjects were given 0.4 mL of the CelWel supplement 4-6 times per day for 14 days. The severity of long COVID symptoms was assessed using the Fatigue Severity Scale Questionnaire (FSSQ) and the Yorkshire COVID-19 Rehabilitation Screening Test (C19-YRS) before and after treatment. In addition, plasma levels of pro-inflammatory cytokines and chemokines and the post-acute sequelae score of COVID-19 (PASC) were also assessed. Safety parameters such as adverse events, haematology, and serum biochemistry were also evaluated.

Results: Results showed that all COVID-19 survivors had higher FSSQ, C19-YRS, and PASC scores along with elevated plasma levels of pro-inflammatory cytokines and chemokines before treatment. CelWel consumed for 14 days significantly reduced FSSQ and C19-YRS scores and plasma cytokine and chemokine levels. Furthermore, with CelWel treatment, a multiplex based PASC score of cytokines showed a decreasing trend in 11 subjects. All laboratory safety parameters remained within the normal range, and no adverse events were reported during the study period.

Conclusion: These findings suggest that the CelWel supplement is a viable and safe option for reducing the severity of symptoms in patients with Long COVID.

Key Words: CelWel; Fatigue; Long COVID; Cytokines; Clinical trial

INTRODUCTION

The SARS-CoV-2 coronavirus (COVID-19) has affected more than 750 million people around the world to date [1]. Acute manifestations of COVID-19 widely affect the pulmonary, cardiovascular, neurological, hematological, and gastrointestinal systems [2]. Its long-term impacts are not well known, but the large numbers of COVID-19 survivors represent post-COVID-19 sequelae [3]. The etiology of post-COVID symptoms

is unknown; however, several factors have been proposed to be involved, including comorbidities, post-critical illness, and immune and inflammatory damage [4]. The most common signs and symptoms reported by COVID survivors include fatigue, cognitive decline, dyspnea, anxiety, weakness, shortness of breath, depression, palpitations, arthralgia, chest and muscle pain which impair their mobility, daily activities, and health related quality of life (HR-QoL) [5-8]. Fatigue is one of the most common and severe post-COVID symptoms among

Received:	29-March-2023	Manuscript No:	ipdehc-23-16040
Editor assigned:	31-March-2023	PreQC No:	ipdehc-23-16040 (PQ)
Reviewed:	14-April-2023	QC No:	ipdehc-23-16040
Revised:	19-April-2023	Manuscript No:	ipdehc-23-16040 (R)
Published:	26-April-2023	DOI:	10.21767/2049-5478.20.03.11

Corresponding author Alben Sigamani, Department of Health Sciences, Carmel Research Consultancy, India, E-mail: dralbens@myrescon.com

Citation Sigamani A, Kumar SS, Naik KS (2023) Effects of a Natural Extract on Molecular Signature in Post-Acute Sequelae SARS-CoV-2 (PASC) Infection: A Quasi-Experimental Study. Divers Equal Health Care. 20:11.

Copyright © 2023 Chen Y. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

52-58% of COVID-19 survivors and the symptom has been reported to persist for more than a year after the onset of illness [9,10]. Fatigue is characterized by an ongoing experience of weakness, exhaustion, and increased difficulty doing physical (e.g., walking, climbing stairs, or lifting) or mental work [11]. In post-COVID syndrome, fatigue is frequently accompanied by muscle fatigue and fatiguability, which coincides with markers of inflammation and hypo-perfusion [12].

Prolonged fatigue after COVID-19 shares clinical manifestations with other post-infectious fatigue syndromes as well as myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) [13]. In both situations, there are higher levels of circulating pro-inflammatory cytokines (such as IL-1, IL-4, GM-CSF, IFN- γ , TNF- α , etc.) and CD8+ T cells, which are presumed to be involved in the pathophysiology of these illnesses. In such chronic inflammatory conditions, as well as possibly in prolonged COVID, inflammatory cytokines reduce basal ganglia function and dopamine while increasing neurotransmitter release in the brain, including serotonin, which causes persistent fatigue [14,15]. In patients recovering from West Nile virus infections, pro-inflammatory (e.g., IL-2, IL-6) and antiviral (e.g., IFN- γ) cytokines were found to be elevated [16]. In contrast, the administration of IL-6 caused fatigue and sleep disturbances in healthy patients [17]. A similar mechanism could be at work in long-COVID fatigue. Considering the prominent role of inflammatory cytokines in longstanding COVID-19 symptoms, it was hypothesised that blocking pro-inflammatory cascades might alleviate longstanding COVID-19 symptoms, particularly fatigue.

CelWel is an oral supplement containing polyphenols derived from *Tinospora cordifolia* and *Piper nigrum*, as well as a nano dose of thimerosal. Polyphenols from *T. cordifolia* in CelWel, has proven anti-inflammatory activity; lowering pro-inflammatory cytokine levels (IL-1, TNF- α , IL-6, and IL-17), reduced frequency of IL-17-producing T cells, and lowering production of chemokines (RANTES) [18]. This study aimed to evaluate the efficacy and safety of CelWel as a supplement for reducing Long COVID symptoms, specifically fatigue, and resetting of various inflammatory biomarkers.

METHODS

Study Design, and Setting

This single-arm, non-randomized, open-label prospective study of CelWel was conducted from December 2022 to January 2023, at Government Medical College and Government General Hospital, Srikakulam, Andhra Pradesh, India. The efficacy and safety of the CelWel supplement were evaluated as adjunctive therapy for a maximum of 14 days for COVID-19 survivors with persistent COVID-19 symptoms. The study was carried out in accordance with the Declaration of Helsinki, the Indian Council of Medical Research (ICMR) ethical guidelines for biomedical research, and ICH guidelines for good clinical practice (GCP). This study protocol was approved by the institutional medical ethical committee (Protocol no.: CW/C-19/1122, Version 1.0) and was prospectively registered with the Clinical Trials Registry-India (ID: CTRI/2022/12/048144) dated 15 December, 2022. All potential participants were voluntarily recruited, and informed consent was obtained prior to the initiation of any study-related procedures.

Participants

Following permission from Institutional Ethics committee, the electronic database of the Government Medical College and Government General Hospital, Srikakulam, Andhra Pradesh, was used to identify potential participants infected with SARS-CoV-2. Male and female patients over the age of 18 who had been diagnosed with COVID-19 by RT-PCR and had persistent mild to moderate post-COVID symptoms lasting 4 to 12 weeks since disease onset were recruited. Patients were excluded if they were asymptomatic, had severe systemic diseases, uncontrolled comorbid conditions, vascular disorders, or active cancer, were hospitalized or required oxygen support, or had received antiviral medication in the last 36 hours. Potential participants were screened for eligibility, and demographic data was collected at the outset.

Interventions

CelWel is an oral liquid supplement, and each 100 mL contains *T. cordifolia* extract (200 mg), *P. nigrum* extract (2.5 mg), and thimerosal (0.05 mg) as active ingredients. Subjects were instructed to take 0.4 mL of the CelWel supplement 4-6 times daily, 5 minutes before eating or drinking. After 14 days of treatment, the efficacy and safety were evaluated.

Primary Outcomes

The primary outcomes were improvements in chronic fatigue and other symptoms associated with long COVID patients as measured by fatigue severity scale questionnaires (FSSQ) [19] and a modified version of the COVID-19 Yorkshire Rehabilitation Screening (C19-YRS) checklist [20]. Fatigue severity was determined using self-administered FSSQ questionnaires (9 items). Each item was scored on a 7-point scale, with 1 representing strongly disagree and 7 representing strongly agree. Higher FSS scores for each item indicate more severe fatigue, and if an FSSQ score ≥ 4 indicates clinically significant fatigue [19]. The final score was the average of the 9 items. Prolonged COVID symptoms were assessed using a modified C19-YRS tool consisting of 18 items rated on an 11-point rating scale as 3=mild symptom, 3-5=moderate symptom, and 6-10=severe symptom. The C19-YRS is well validated tool that correlates well with exercise tolerance, lower limb strength and mobility, anxiety, depression and HR-QoL [8]. The C19-YRS checklist analyzed symptoms related to body functions and structures (12 items), activity limitations (5 items), and personal factors (1 item). After 14 days of treatment, the FSSQ and C19-YRS scores were evaluated.

Secondary Outcomes

Secondary outcomes of this study included changes in plasma levels of inflammatory cytokines (IL-2 and IFN- γ) and chemokines (CCL4 (MIP-1 β)). IL-2, IFN- γ , and CCL4 (MIP-1 β) levels in plasma were measured using multiplex bead-based flow cytometry (IncellKINE, IncellDx, Inc.) [21]. The assay was carried out in a 96-well microplate, and samples were analysed using a CytoFlex LX 3-laser flow cytometer (Beckman Coulter Life Sciences, USA) and Kaluza Analysis Software (Beckman-Coulter, Miami, FL). Changes in laboratory parameters, such as the complete blood count (CBC), ESR, renal function, and liver function parameters from baseline, were evaluated as part of the safety study.

Statistical Analysis

A per-protocol (PP) population analysis was used to assess efficacy. Demographic data were presented as mean, percentage, and standard deviation (SD). Efficacy parameters, such as FSSQ and C19-YRS scores, were compared between baseline data and final follow-up data using the Wilcoxon signed-rank test. To compare the mean change in plasma IL-2, IFN- γ , and CCL4 (MIP-1 β) levels from baseline to the final visit, the Wilcoxon signed-rank test was used. Unless otherwise stated, all hypotheses were tested at a significance level of 0.05. All statistical analysis was carried out using SPSS software version 10.0.

RESULTS

Demographic and Baseline Characteristics

A total of 15 participants were included in the study, of whom 9

Table 1: FSSQ scores after 14 days of treatment with CelWel supplement.

FSSQ	Day 1, N=15 FSSQ score (mean $\pm$ SD)	Day 14, N=15 FSSQ score (mean $\pm$ SD)	Z-Score	P-value
Motivation lower when fatigued	4.80 $\pm$ 1.61	4.00 $\pm$ 0.76	-2.164	0.030*
Exercise brings on my fatigue	5.00 $\pm$ 1.07	3.93 $\pm$ 0.83	-2.739	-2.739
I am easily fatigued	4.67 $\pm$ 1.35	3.80 $\pm$ 0.86	-2.124	0.034*
Fatigue interferes with my physical functioning	4.80 $\pm$ 1.01	3.73 $\pm$ 0.80	-3.066	0.002**
Fatigue causes frequent problems for me	4.80 $\pm$ 0.78	3.33 $\pm$ 0.82	-3.115	0.002**
My fatigue prevents sustained physical functioning	4.87 $\pm$ 0.92	3.60 $\pm$ 0.83	-2.994	0.003**
Fatigue interferes with carrying certain out certain duties and responsibilities	4.93 $\pm$ 0.96	3.67 $\pm$ 0.98	-3.079	0.002**
Fatigue is among my 3 most disabling symptoms	4.60 $\pm$ 0.83	3.27 $\pm$ 0.88	-3.115	0.002**
Fatigue interferes with work, family & social life	4.87 $\pm$ 0.92	3.27 $\pm$ 1.03	-3.247	0.001***
Total fatigue scores	43.30 $\pm$ 7.77	32.30 $\pm$ 5.35	-3.295	0.001***

Assessment of long COVID symptoms by C19-YRS checklist: The severity of long COVID symptoms was assessed by the C19-YRS checklist at days 1 and 14. Among the Body Functions and Structures items, the intensity of the breathlessness on climbing stairs, fatigue, continence, cognition, anxiety, and depression at day 1 were the prominent symptoms compared to other symptoms, with C19-YRS reporting intensities of 5.43 $\pm$ 1.22, 6.08 $\pm$ 1.04, 6.0 $\pm$ 0.78, 5.17 $\pm$ 1.27, 6.00 $\pm$ 0.78, 5.40 $\pm$ 0.91 and 5.27 $\pm$ 1.10, respectively (Table 2). After 14 days of treatment,

(60%) were men and 6 (40%) were women. The subjects average is 47.0 $\pm$ 10.58 years. The mean height, weight, and BMI of the participants are 161 $\pm$ 8.16 cm, 61.60 $\pm$ 7.49 kg, and 23.71 $\pm$ 1.17 kg/m², respectively.

Primary Outcomes

Assessment of fatigue: The fatigue severity in COVID-19 survivors was assessed using the FSSQ on days 1 and 14. The results revealed that patients consistently had higher FSSQ scores (i.e., >4) for each item as well as a higher mean FSSQ score at day 1. The FSSQ score of each item was significantly reduced after 14 days of CelWel treatment, and the changes were statistically significant (Table 1). The mean FSSQ score also decreased significantly ($p < 0.001$) from day 1 (43.30 $\pm$ 7.77) to day 14 (32.30 $\pm$ 5.35).

all these symptoms were present but significantly reduced. Improvements in fatigue, continence, cognition, anxiety, and depression symptoms were found to be statistically significant. Among the activities, a statistically significant improvement in mobility, personal care, usual activities, and social role were observed after 14 days of treatment compared to day 1. The global health perception was also statistically significantly reduced from 3.47 $\pm$ 1.51 (day 1) to 2.33 $\pm$ 1.50 (day 14).

Table 2: Intensity change in long COVID symptoms assessed by C19-YRS.

Activities	Day 1, N=15 C19-YRS score (mean $\pm$ SD)	Day 14, N=15 C19-YRS score (mean $\pm$ SD)	Z-Score	P-value
Body Functions and Structures				
Breathlessness at rest	4.13 $\pm$ 1.96	4.27 $\pm$ 2.02	-0.816	0.414
Breathlessness dressing	2.80 $\pm$ 1.01	2.87 $\pm$ 1.06	-1.0	0.317
Breathlessness stairs	5.43 $\pm$ 1.22	5.25 $\pm$ 1.67	-1.0	0.317
Cough/throat sensitivity/ voice	3.40 $\pm$ 1.34	3.33 $\pm$ 0.58	-0.447	0.655
Swallowing	2.50 $\pm$ 0.71	2.80 $\pm$ 2.39	na	na
Fatigue	6.08 $\pm$ 1.04	4.85 $\pm$ 1.41	-2.858	0.004**

Continence	6.00 ± 0.78	4.43 ± 1.22	-3.236	0.001***
Cognition	5.17 ± 1.27	4.25 ± 1.71	-2.373	0.018*
Pain/discomfort	1.50 ± 0.58	2.50 ± 1.69	-0.447	0.655
Anxiety	5.40 ± 0.91	3.87 ± 1.55	-3.108	0.002**
Depression	5.27 ± 1.10	3.29 ± 1.64	-3.108	0.002**
PTSD screen	1.50 ± 0.71	0.50 ± 0.71	-1.0	0.317
Activities				
Communication	1.80 ± 1.30	1.80 ± 1.30	-1.414	0.157
Mobility	5.60 ± 1.50	5.60 ± 1.50	-3.354	0.001***
Personal care	5.14 ± 1.99	5.14 ± 1.99	-2.831	0.005**
Usual activities	5.20 ± 1.78	5.20 ± 1.78	-3.105	0.002**
Social role	5.13 ± 1.46	5.13 ± 1.46	-3.471	0.001***
Personal Factors				
Global health	3.47 ± 1.51	2.33 ± 1.50	-2.358	0.018**

Secondary Outcomes; Assessment of inflammatory cytokines and chemokines: Sum-marizes the plasma levels of IL-2, IFN- γ , and CCL4 (MIP-1 β). Plasma levels of IL-2, IFN- γ , and CCL4 (MIP-1 β) were measured on days 1 and 14 (**Table 3**). IL-2 levels increased slight-ly from 4.74 ± 2.45 (day 1) to 5.01 ± 2.90 (day 14), whereas IFN- γ and CCL4 (MIP-1 β) levels decreased from 37.58 ± 86.27 to 18.23 ± 22.61 and 1527.07 ± 2185.50 to 1883.81 ± 3834.16, respectively. However, the changes were not statistically sig-nificant. We calculated the post-acute sequelae of COVID-19 (PASC) disease score based on the findings using the Patterson et al. [21] proposed formula: $S1 = (IFN-\gamma + IL-2) / CCL4 (MIP-1\beta)$

with optimized threshold for $S1 = 0.5$. At day 1, all patients had a higher PASC score (>0.5), and after 14 days of treatment, there was a trend in the reduction of PASC score among the 11 participants, but 4 patients showed a reversal of the trend in PASC score (**Table 4**).

Safety: The safety parameters included CBC, ESR, and liver and kidney function parameters that were evaluated before and after 14 days of treatment. On day 14, analysis of safety parameters showed no significant differences compared to day 1, and all parameters were within the normal range (**Tables 5 and 6**). There were no adverse or serious reactions reported during the treatment period.

Table 3: Plasma levels of cytokines/chemokines before and after treatment

Cytokine/ Chemokine	Day 1, N=15 Concentration (mean ± SD)	Day 1, N=15 Concentration (mean ± SD)	P-value
IL-2	4.74 ± 2.45	5.01 ± 2.90	0.865
IFN- γ	37.58 ± 86.27	18.23 ± 22.61	0.191
CCL4 (MIP-1 β)	1883.81 ± 3834.16	1527.07 ± 2185.50	0.955

Table 4: PASC Score among the participants before and after treatment

Patients	PASC Score (Day 1)	PASC Score (Day 14)	Score diff.
P01	8.80	2.79	-6.01
P02	7.40	9.31	1.91
P03	4.31	2.77	-1.54
P04	3.82	23.63	19.81
P05	345.85	72.36	-273.49
P06	2.69	2.04	-0.66
P07	11.22	3.83	-7.38
P08	25.95	3.16	3.16
P09	11.24	48.82	48.82
P10	23.94	17.77	17.77
P11	19.50	11.09	11.09
P12	50.38	4.32	-46.06
P13	3.03	55.90	52.87
P14	29.46	2.32	-27.14
P15	18.75	16.78	-1.97

Table 5: CBC parameters and ESR before and after treatment

Cytokine/ Chemokine	Day 1, N=15 Concentration (mean ± SD)	Day 14, N=15 Concentration (mean ± SD)	P-value
RBC (Million/c. mm)	3.76 ± 0.3	3.81 ± 0.27	1.0
Hb (gm/dl)	12.27 ± 1.59	12.39 ± 1.51	0.102
WBC (cells/ c.mm)	8340 ± 1000.67	8373.33 ± 1230.86	0.683

Neutrophils (%)	60.47 ± 2.33	60.73 ± 1.53	0.509
Lymphocytes (%)	28.53 ± 1.6	29.73 ± 0.8	0.017
Monocytes (%)	2.0 ± 1.41	4.27 ± 1.79	0.006
Eosinophils (%)	8.67 ± 2.09	5.07 ± 1.87	0.001
Basophils (%)	0.4 ± 0.83	0.2 ± 0.41	0.527
Platelet count (lakhs)	2.32 ± 0.35	2.35 ± 0.32	0.317
ESR (mm 1st hour)	8.67 ± 4.37	4.87 ± 3.27	0.001

Table 6: Serum biochemical parameters before and after treatment

Cytokine/ Chemokine	Day 1, N=15 Concentration (mean ± SD)	Day 14, N=15 Concentration (mean ± SD)	P-value
RBS (mg/dl)	110.53 ± 14.6	107.47 ± 10.91	0.317
BUN (mg/dl)	9.07 ± 2.09	9.13 ± 2.0	0.785
Sr. uric acid (mg/dl)	2.47 ± 1.06	2.47 ± 0.99	1.0
Sr. creatinine (mg/dl)	0.72 ± 0.12	0.72 ± 0.11	1.0
Sr. albumin (gm%)	3.71 ± 0.46	3.68 ± 0.51	1.0
Bilirubin (mg/dl)	0.63 ± 0.13	0.62 ± 0.14	0.317
ALT (IU/L)	39.53 ± 9.53	38.07 ± 8.46	0.012
AST (IU/L)	37.2 ± 11.07	36.47 ± 9.85	0.235

DISCUSSION

Long-term COVID syndrome (the post-acute sequelae of SARS-CoV2 infection) is a growing public health problem. There is increasing evidence that a significant proportion of COVID-19 survivors develop a chronic and debilitating set of multisystem signs and symptoms [22]. It has a negative impact on society, the economy, and patient's quality of life, physical functioning, work participation, care dependency, and activities of daily living [23]. Long COVID has been associated with over 60 physical and psychological symptoms [24]. Profound fatigue is one of the most commonly reported symptoms in post-acute COVID-19 patients, with a prevalence of 52% within 3 weeks of hospital discharge [25]. The persistence of fatigue following an acute COVID infection is also associated with episodes of overwhelming exhaustion, depressive symptoms, poor sleep quality, memory impairment and low energy levels, all of which influence quality of life [26]. There is currently no specific pharmacotherapy available for the treatment of long COVID. Recently, efforts have been made to understand the cause, epidemiological burden, and potential biomarkers of long COVID [27].

This study aimed to evaluate the efficacy of the CelWel product in improving long COVID symptoms, particularly persistent fatigue in COVID-19 survivors. The findings revealed that all patients experienced clinically significant fatigue prior to CelWel treatment, as evidenced by a higher FSSQ score (≥ 4) for each question as well as an overall FSSQ score (>40). After 14 days of treatment, CelWel significantly reduced FSSQ scores. The C19-YRS test was also used in this study to assess the effect of CelWel on post-COVID symptoms. Results showed that CelWel significantly improved some symptoms related to the body's functions and structures, including fatigue and limitations in activities. Therefore, CelWel was found to be an effective health supplement to ameliorate long COVID symptoms, including persistent fatigue in COVID-19 survivors. However, its exact mechanism for reducing persistent fatigue and other long COVID symptoms is unknown.

It has recently been suggested that an ongoing low-grade pro-inflammatory response may be linked to the development of post-COVID symptoms [28]. As mentioned in the introduction, long COVID and ME/CFS share some clinical manifestations. ME/CFS pathophysiology has been linked to immune function dysregulation, hyperinflammation, oxidative stress, and autoimmunity [29]. Furthermore, ME/CFS-associated post-infectious fatigue syndrome has been linked to other acute viral and bacterial infections such as severe acute respiratory syndrome (SARS), Middle East respiratory syndrome (MERS), Epstein-Barr virus (EBV), human parvovirus (HPV)-B19, and Q fever [23,30]. Moreover, neuro immunological involvement is a crucial precondition for long-term fatigue [31,32]. IFN- α and other cytokines of the innate immune response have been shown to promote behavioural changes in medically ill patients, including fatigue-related symptoms, by acting on the basal ganglia nuclei [33]. Evidence from studies with neuro immunological disorders suggests strong connections between structural brain abnormalities and fatigue severity, such as changes in the basal ganglia [34,35] and from to parietal white matter [36] associated with fatigue in multiple sclerosis. New

structural images of the brain in patients with post-COVID syndrome revealed abnormalities in the thalamus and basal ganglia, which are associated with post-COVID fatigue [26].

With these similarities between Long-COVID and ME/CFS, as well as the neuro-immunological disorders associated with fatigue, it is plausible that SARS-CoV-2 infection can trigger PASC symptoms, particularly fatigue, by inducing an inflammatory state and a dysregulated immune response. We measured plasma levels of IL-2, IFN- γ , and CCL4 (MIP-1 β) to see if CelWel treatment affected patients' immunogenic profiles. CelWel had no effect on the pro-inflammatory cytokine IL-2, but it did down-regulate IFN- γ and CCL4 (MIP-1 β). In addition to that, after CelWel treatment, the PASC score decreased in 11 patients, indicating a reduction of the inflammatory state by preventing the production of pro-inflammatory cytokines in patients with prolonged COVID-19. However, four patients had the opposite trend in PASC score and would have required a large or very prolonged dose of CelWel to improve their PASC score. During the study period, no patients reported adverse events or abnormal clinical findings in laboratory parameters.

The limitations of the present study are the inability to measure other inflammatory cytokines, the non-randomized, small sample size, and the fact that the study was conducted in a single center. A future study should require comparing the effects of CelWel with placebos or standard of care on a large cohort for an extended period.

CONCLUSION

The results of this study indicate that the CelWel supplement has the potential to be an effective and safe treatment for Long-COVID. CelWel shows promise to reverse PASC symptoms by resetting the immune system of COVID survivors. However, the underlying mechanisms of action of CelWel need to be established through a large-scale trial.

AUTHOR CONTRIBUTIONS

Dr. Alben Sigamani, study design and manuscript editing; Dr. K. Sunil Naik, clinical investigation and manuscript editing; Sangeetha Sampath Kumar, manuscript preparation and literature support.

DATA AVAILABILITY

The corresponding author can provide derived data to support the findings of this study upon request.

ACKNOWLEDGEMENTS

We would like to acknowledge the management of the Government Medical College and Government General Hospital (Old RIMS), Srikakulam, for their assistance for this study. We would also like to thank the nursing staff, and support staff of the Hospital for their assistance throughout this study.

REFERENCES

1. World Health Organization. COVID-19 Dashboard, 2022.
2. Zheng KI, Feng G, Liu WY, Targher G, Byrne CD, et al. (2021)

- Extra pulmonary complications of COVID-19: A multisystem disease? *J Med Virol* 93:323-335.
3. Sykes DL, Holdsworth L, Jawad N, Gunasekera P, Morice AH, et al. (2021) Post-COVID-19 symptom burden: What is long-COVID and how should we manage it? *Lung* 199:113-119.
 4. Verveen A, Wynberg E, Willigen HDG, Boyd A, Jong MD, et al. (2022) Recovered study group. Severe fatigue in the first year following SARS-CoV-2 infection: A prospective cohort study. *Open Forum Infect Dis* 9(5):ofac127.
 5. Daugherty SE, Guo Y, Heath K, Dasmariñas MC, Jubilo KG, et al. (2021) Risk of clinical sequelae after the acute phase of SARS-CoV-2 infection: Retrospective cohort study. *BMJ* 373:n1098.
 6. Hansen KS, Mogensen TH, Agergaard J, Christensen B, Ostergaard L, et al. (2023) High-dose coenzyme Q10 therapy versus placebo in patients with post COVID-19 condition: A randomized phase 2, crossover trial. *Lancet Reg Health Eur* 24:100539.
 7. Kunoor A, Surendran D, Hari H, Viswan V, Harikrishnan K, et al. (2022) Impact of early pulmonary rehabilitation in postacute COVID Disease: A single-center experience from India. A quasi-experimental study. *Indian J Public Health*. 66:S51-S55.
 8. Straudi S, Manfredini F, Baroni A, Milani G, Fregna G, et al. (2022) Construct validity and responsiveness of the COVID-19 yorkshire rehabilitation scale (C19-YRS) in a cohort of Italian hospitalized COVID-19 patients. *Int J Environ Res Public Health*. 19:6696.
 9. Cuadrado ML, Florencio LL, Santiago OR, Martínez CM, Trigueros LJ, et al. (2022) Fatigue and dyspnoea as main persistent post-COVID-19 symptoms in previously hospitalized patients: Related functional limitations and disability. *Respiration* 101:132-141.
 10. Yong SJ (2021) Long COVID or post-COVID-19 syndrome: Putative pathophysiology, risk factors, and treatments. *Infect Dis (Lond)* 53:737-754.
 11. Paniagua CJ, Arribas MJ, Calero JA, Vidal MI, Peñas FC, et al. (2022) A tele-health primary care rehabilitation program improves self-perceived exertion in COVID-19 survivors experiencing Post-COVID fatigue and dyspnea: A quasi-experimental study. *PLoS One* 17:e0271802.
 12. Kedor C, Freitag H, Arndt ML, Wittke K, Hanitsch LG, et al. (2022) A prospective observational study of post-COVID-19 chronic fatigue syndrome following the first pandemic wave in Germany and biomarkers associated with symptom severity. *Nat Commun* 13:5104.
 13. Komaroff AL, Lipkin WI (2021) Insights from myalgic encephalomyelitis/chronic fatigue syndrome may help unravel the pathogenesis of postacute COVID-19 syndrome. *Trends Mol Med* 27:895-906.
 14. Jennifer C. Felger AHM (2012) Cytokine effects on the Basal Ganglia and dopamine function: The subcortical source of inflammatory malaise. *Front Neuroendocrinol* 33:315.
 15. Cordeiro LMS, Rabelo PCR, Moraes MM, Coimbra CC, Wanner SP, et al. (2017) Physical exercise-induced fatigue: The role of serotonergic and dopaminergic systems. *Braz J Med Biol Res* 50:e6432.
 16. Islam MF, Cotler J, Jason LA (2020) Post-viral fatigue and COVID-19: Lessons from past epidemics. *Fatigue: Biomed Health Behav* 8:61-69.
 17. Späth-Schwalbe E, Hansen K, Schmidt F, Schrezenmeier H, Marshall L, et al. (1998) Acute effects of recombinant human interleukin-6 on endocrine and central nervous sleep functions in healthy men. *J Clin Endocrinol Metab* 83:1573-1579.
 18. Sannegowda KM, Venkatesha SH, Moudgil KD (2015) *Tinospora cordifolia* inhibits autoimmune arthritis by regulating key immune mediators of inflammation and bone damage. *Int J Immunopathol Pharmacol* 28:521-531.
 19. Valko PO, Bassetti CL, Bloch KE, Held U, Baumann CR, et al. (2008) Validation of the fatigue severity scale in a Swiss cohort. *Sleep* 31:1601-1607.
 20. Sivan M, Halpin S, Gee J (2020) Assessing long-term rehabilitation needs in COVID-19 survivors using a telephone screening tool (C19-YRS tool). *Adv Clin Neurosci Rehabil* 19:14-17.
 21. Patterson BK, Coto GJ, Yogendra R, Francisco EB, Long E, et al. (2021) Immune based prediction of COVID-19 severity and chronicity decoded using machine learning. *Front Immunol* 12:700782.
 22. Hanson SW, Abbafati C, Blyuss O (2022) Global burden of disease long COVID collaborators. Estimated global proportions of individuals with persistent fatigue, cognitive, and respiratory symptom clusters following symptomatic COVID-19 in 2020 and 2021. *JAMA* 328:1604-1615.
 23. Herck VM, Goërtz YMJ, Wilke HS, Machado FVC, Delbressine JM, et al. (2021) Severe fatigue in long COVID: Web-based quantitative follow up study in members of online long COVID support groups. *J Med Internet Res* 23:e30274.
 24. Michelen M, Manoharan L, Elkheir N, Cheng V, Dagens A, et al. (2021) Characterising long COVID: A living systematic review. *BMJ Glob Health* 6:e005427.
 25. Leon LS, Ostrosky ST, Perelman C, Rebolledo PA (2021) More than 50 long-term effects of COVID-19: A systematic review and meta-analysis. *Sci Rep* 11:16144.
 26. Heine J, Hartung TJ, Rekers S, Chien C, Boesl F, et al. (2023) Structural brain changes in patients with post-COVID fatigue: A prospective observational study. *E Clin Med* 58:101874.
 27. Koc HC, Xiao J, Liu W, Li Y, Chen G, et al. (2022) Long COVID and its management. *Int J Biol Sci* 18:4768-4780.
 28. Afrin LB, Weinstock LB, Molderings GJ (2020) COVID-19 hyper inflammation and post-COVID-19 illness may be rooted in mast cell activation syndrome. *Int J Infect Dis* 100:327-332.
 29. Rivera MC, Mastronardi C, Aldana CT, Burgos AM, Lidbury

- BA, et al. (2019) Myalgic encephalomyelitis/chronic fatigue syndrome: A Compr Rev. Diagn 9:91.
30. Hernandez OD, Shoenfeld Y (2009) Infection, vaccination, and autoantibodies in chronic fatigue syndrome, cause or coincidence? *Ann NY Acad Sci* 1173:600-609.
 31. Frontera JA, Yang D, Lewis A, Patel P, Medicherla C, et al. (2021) A prospective study of long-term outcomes among hospitalized COVID-19 patients with and without neurological complications. *J Neurol Sci* 426:117486.
 32. Hartung TJ, Neumann C, Bahmer T, Endres M (2022) Fatigue and cognitive impairment after COVID-19. *E Clin Med* 53:101651.
 33. Capuron L, Pagnoni G, Demetrashvili MF, Lawson DH, Fornwalt FB, et al. (2007) Basal Ganglia hyper metabolism and symptoms of fatigue during interferon- α therapy. *Neuropsychol* 32: 2384-2392.
 34. Jaeger S, Paul F, Scheel M, Brandt A, Heine J, et al. (2019) Multiple sclerosis-related fatigue: Altered resting-state functional connectivity of the ventral striatum and dorso-lateral prefrontal cortex. *Mult Scler J* 25:554-564.
 35. Fleischer V, Ciolac D, Grothe M, Strauss S, Galindo LS, et al. (2022) German competence network multiple sclerosis (KKNMS). Subcortical volumes as early predictors of fatigue in multiple sclerosis. *Ann Neurol* 91:192-202.
 36. Yarraguntla K, Bao F, Mikol LS, Razmjou S, Martinez SC, et al. (2019) Characterizing fatigue-related white matter changes in MS: A proton magnetic resonance spectroscopy study. *Brain Sci.* 9(5):122.